


bioRxiv preprint doi: <https://doi.org/10.1101/052480>; this version posted September 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

FIGURE 1A

Viability - MCF7



Viability - SKBR3




FIGURE 1C

Viability - HTB 132





FIGURE 1D

Effect of reovirus on MCF7 viability



Reovirus DNA Fragmentation MCF-7




FIGURE 2B

Reovirus DNA Fragmentation SKBR3




FIGURE 2C

Reovirus DNA Fragmentation HTB 132




FIGURE 2D

Apoptosis (Annexin V-/7AAD)




FIGURE 2E

Apoptosis (APO 2.7) - MCF7 cells




FIGURE 2F

Apoptosis (APO 2.7) - HTB 132 cells




FIGURE 2G

Apoptosis (APO 2.7) - SKBR3 cells





FIGURE 3A


Effect of reovirus on CD34+ cells




Hours post virus addition

FIGURE 3B


Effect of reovirus on long- term stem cell culture



Purging apheresis product of contaminating MCF-7 cells



Purging apheresis product of contaminating HTB-132 cells



Purging apheresis product of contaminating SKBR3 cells

